

Research Article

Published Date:-2023-12-19 15:00:04

[Success, Survival and Prognostic Factors in Implant Prosthesis: Experimental Study](#)

The primary objective of this study was to detect the success and short-term survival rate of dental implant prosthetic therapy.

The valuation of a possible relationship between the general and local clinical conditions of the patients (presence of risk factors and type of dental implant-supported prosthesis) and the satisfaction perceived by the patient, with success and survival of implant devices was investigated.

The sample trial consisted of 23 patients, for a total of 50 dental implants supporting a prosthetic therapy.

Preliminarily, an analysis of averages and frequencies of the anamnestic data was presented, as numbers and percentages. The implant success rate was calculated by assessing whether the implants fulfilled the success criteria defined by the Pisa Consensus Conference. The ANOVA test was used to check whether there was an association between the success of the implant device, the anamnestic data collected, and the type of prosthesis supported by the implants. Finally, the survival rate was calculated using the Kaplan - Meier method.

The 2-year success rate of 50 implants was assessed at 98%. The 2-year survival rate was 97%. Finally, sex, age, compensated diabetes, a smaller and equal number of cigarettes per day (10 per day), the BOP, and the type of titanium dental implants supported by prosthesis, do not determine a change in success and are therefore not adequate parameters to predict the outcome of implant success. Following the results obtained, it is appropriate to continue the research by expanding the clinical observation times in order to obtain more solid scientific and clinical evidence.

Research Article

Published Date:-2023-08-08 14:51:35

[Knowledge, Attitude, and Practices of Parents toward \(Infant & Child\) Oral Health in Family Medicine Center at PSMCC, Riyadh](#)

Background: Early childhood caries may develop as soon as teeth erupt. Promoting oral health care programs is very important and helps to reduce dental caries in young children. The aim of this study was to assess the infant/child oral health (IOH) related knowledge, attitudes, and practices (KAP) of parents in Riyadh, Saudi Arabia. Materials and Methods: A cross-sectional descriptive study was conducted among 327 parents. Visiting the family medicine center at prince sultan military medical city, Riyadh Saudi Arabia. A 39-item questionnaire covering socio-demographic characteristics and questions pertaining to KAP regarding OH care will be used to collect the data. Descriptive statistics, Student's t-test, one-way analysis of variance, and Scheffe's test will use for the statistical analysis ($p < 0.05$).

Results: The result showed that the majority of the parents had good knowledge regarding OH, knowledge of cleaning (92.4%), and knowledge of the amount of sugar (88.1%). the parent's age group (30 to 39) years ($n = 147$) reported the highest mean (knowledge, attitudes, and practice) scores among all other age groups with a knowledge mean score of (6.80 ± 1.73), an attitude mean score of (8.86 ± 1.37), and a practice mean score of (5.14 ± 1.86).

Female parents showed significantly higher mean knowledge, attitude, and practice scores than male parents. In addition, the middle-income level parents' group ($n = 295$) reported higher knowledge mean score compared to the low-income parents' group ($n = 15$) with a mean difference of $d = 1.15$, $p = .041$.

Conclusion: Parents' knowledge about maintaining oral and dental health care for infants/children was inadequate. Essentially, medical professionals are the initial ones to interact with expecting and new moms. Therefore, need to raise parents' awareness about oral and dental health for infants/children, through the development and implementation of long-term education and promotions programs.
